Dr. Curtis Noel's Protocol for: ANTERIOR BANKART REPAIR		CRYSTAL CLINIC ORTHOPAEDIC CENTER Apartnership with Summa Health System	
Sling Use:		Sling for 8 weeks ***Avoid the "throwing position" for 3 months***	
Phase I: (Passive)	Week 1-3	 Pendulums to warm-up Passive Range of Motion Supine External Rotation – 0 degrees Supine Forward Elevation - 90 °degrees No Internal Rotation 	
	Week 4	 Supine External Rotation – 30 ° degrees Supine Forward Elevation – Full Internal Rotation to belt line 	
Phase II: (Active/Assistive)	Week 5	 Pendulums to warm-up Active Assistive Range of Motion with Passive Stretch to prescribed limits. Supine-Seated External Rotation - gradually increase to full by 12 weeks. No terminal stretching with ER. Supine-Seated Forward Elevation - Full Internal Rotation - Full 	
Phase III: (Resisted)	Week 6	 Pendulums to warm up and continue with phase 2 External and Internal Rotation Standing forward punch Seated rows Bicep Curls PRE's for Periscapular strengthening 	
Weight Training:	Week 12	Keep Min	id anterior capsular stress o hands within eyesight, Keep elbows bent imize overhead activities Military press, pulldown behind head, or wide grip ch)
	Immediately		Computer, eating, holding a book, typing, writing
Return to activities:	8 weeks		Golf (chip and putt ONLY)
	12 weeks		Tennis (no overhead)
	4 months		Contact Sports
	6 months		Throwing program